

LA-UR-18-24798

Approved for public release; distribution is unlimited.

Title: Antarctic Ice shelf-ocean interactions in high-resolution, global

simulations using the Energy Exascale Earth System Model (E3SM)

Author(s): Asay-Davis, Xylar Storm

Comeau, Darin Scott Hoffman, Matthew James Maltrud, Mathew Einar Petersen, Mark Roger Price, Stephen F. Dr Van Roekel, Luke Veneziani, Carmela Wolfram, Phillip Justin Jr.

Intended for: European Seminar on Computing, 2018-06-04/2018-06-08 (Pilsen, Czech

Republic)

Issued: 2018-06-04








Antarctic Ice shelf-ocean interactions in high-resolution, global simulations using the Energy Exascale Earth System Model (E3SM)

Xylar Asay-Davis, Darin Comeau, Matthew Hoffman, Mathew Maltrud, Mark Petersen, Stephen Price, Luke Van Roekel, Milena Veneziani, Phillip Wolfram

Los Alamos National Laboratory





Outline

- E3SM model and project
- Ice shelf/ocean interactions in E3SM
- Land ice/ocean coupling in E3SM
- Summary





Energy Exascale Earth System Model (E3SM)*

The U.S. Department of Energy (DOE) *E3SM* project aims to meet DOE and U.S. government agency needs for state-of-the-science Earth system modeling.

- Phase 1: July 2014 June 2018
- Public release: April 2018
- Phase 2: July 2018 June 2021

Science focus areas:

- Water Cycle
- Biogeochemistry
- Cryosphere:

How will the atmosphere, ocean, and sea-ice systems mediate sources of sea-level rise from the Antarctic ice sheet over the next 30 years?

N. Atlantic hurricane in high-resolution configuration (color=SST)

^{*} formerly the Accelerated Climate Model for Energy (ACME)





E3SM Version 1.0 Release

Public release of v1 code and simulations ~6 weeks ago

- Project homepage (https://e3sm.org)
- Code available on Github (https://github.com/E3SM-Project/E3SM)
- Simulations (limited DECK) (https://e3sm.org/data/get-e3sm-data)

Highlights:

- open development
- variable resolution (all components)
- new ocean, sea ice, land ice models
- new var. res. analysis framework
- ocean circulation beneath ice shelves (currently unverified; unsupported)






Cryosphere Simulations (ongoing)

Identical low (EC60to30km) & medium (RRS30to10km) resolution configurations, with & without Antarctic ice shelves:

- Prescribed atmospheric forcing (multiple CORE-IAF cycles)
- Coupled atmosphere/land/ocean/ sea-ice pre-industrial (several cen
- Coupled atmosphere/land/ocean/ sea-ice historical (1850-present)




ocean transport in global, med. resolution simulation





Cryosphere Simulations: Goals

- Demonstrate ice shelves "do no harm" to global simulations
- Demonstrate sub-ice shelf melt rates and other relevant ice-sheet-proximal climate characteristics plausible relative to observations
- Explore sensitivity to (1) presence/ absence of shelves, (2) forcing and coupling, (3) resolution
- Release of CORE-IAF, pre-industrial equilibrium, and historical simulations by fall 2018




ocean transport in global, med. resolution simulation





Ice shelf/ocean interactions

- Ice shelves: floating extensions of ice sheets and glaciers
- Ocean waters flow into cavities below ice shelves
- Temperature of inflowing water determines the amount of melting (nonlinear effect)




From Hanna et al. (2013)





Observed Bottom Temperature and Melt Rates




From Schmidtko et al. (2014)

From Rignot et al. (2013)






Ice shelves in E3SM








Ocean temp. sensitivity to Ice Shelves in E3SM








Ocean salinity sensitivity to Ice Shelves in E3SM








Melt rate sensitivity to resolution in E3SM








Melt rate sensitivity to resolution in E3SM













Melt rate sensitivity to resolution in E3SM















Seafloor temp. sensitivity to resolution in E3SM








Seafloor temp. sensitivity to resolution in E3SM











Surface velocity sensitivity to resolution in E3SM









Implementation of ice-shelf cavities in E3SM

- Vertical coordinate system:
 - constrained pressure-gradient errors
- Adjustment of sea-surface height:
 - must be consistent with ice pressure
- Parameter study:
 - Best-fit values for uncertain coefficients





Horizontal pressure gradient with tilted coordinates

Horizontal pressure gradient:

$$-\frac{\partial P}{\partial x}\Big|_{z} = -\rho_{s}g\frac{\partial \zeta}{\partial x} - g\int_{z}^{\zeta} \mathcal{J}(\rho, z)ds,$$


• Green's Theorem:

$$\iint\limits_{\mathcal{A}} \mathcal{J}(\rho, z) \ dx \ ds = \oint\limits_{\mathcal{A}} \rho(\mathbf{k}, \mathbf{l}) dl,$$

Nonlinear equation of state (EOS):

$$\rho = \rho(S, \Theta, p)$$

EOS involves the pressure itself



From Shchepetkin and McWilliams (2003)




Horizontal pressure gradient with tilted coordinates

Second-order discretization of

$$\oint \rho(\mathbf{k}, \mathbf{l}) dl,$$

leads to instability when Layers are both:

- thin
- strongly tilted
- Haney Number rx1: a non-dimensional measure of this effect (Haney 1991)




From Shchepetkin and McWilliams (2003)





The Haney Number: rx1 ≤ 5








Consistent sea surface height and ice pressure


- Ice pressure is the weight of the overlying ice shelf
- In equilibrium, balanced by ocean pressure
- Pressure at ice-ocean interface (z_b)
 changes by raising/lowering





Consistent sea surface height and ice pressure

- Difficulties in finding equilibrium z_b:
 - density variation (horizontal and vertical)
 - Various possible equations of state
 - Various possible implementations of horizontal pressure gradient



 E3SM uses iterative adjustment of z_b to minimize dz_b/dt over the first hour of sim.





Parameter Sensitivity Study at 30-km resolution

- 23 runs with 13 uncertain parameters
- Compared melt rates to
 observations (Rignot et al. 2013)
- Found 4 parameters with high sensitivity
- Determined best-fit values to minimize bias

Green's function method of Menemenlis et al. (2005):

 Assume perturbations in model state are linear in perturbed parameter values:

$$y^d = G\eta + \varepsilon$$

• Find parameter values that minimize a cost function:

$$J = \boldsymbol{\varepsilon}^T \boldsymbol{R}^{-1} \boldsymbol{\varepsilon} = \sum_{i} \left(\frac{y_i^0 - x_i}{\sigma_i} \right)^2$$

• Best-fit parameter perturbations are computed as:

$$\boldsymbol{\eta_{fit}} = \left(\boldsymbol{G}^T \boldsymbol{R}^{-1} \boldsymbol{G}\right)^{-1} \boldsymbol{G}^T \boldsymbol{R}^{-1} \boldsymbol{y}^d$$





Parameter Sensitivity Study at 30-km resolution

Parameter Name	Control Value	Perturbed Value	RMS Melt Diff. (Rignot et al. 2013)	Best-fit Value
ctrl	N/A	N/A	3.274	N/A
land_ice_flux_attenuation_coefficient	10	5	3.274	2.56
		15	3.259	
land_ice_flux_boundarylayerthickness	10	5	3.283	0.351
		15	3.268	
land_ice_flux_jenkins_heat_transfer_coefficient land_ice_flux_jenkins_salt_transfer_coefficient	0.011 3.1 × 10 ⁻⁴	0.005 1.4 × 10 ⁻⁴	2.046	0.00295 8.42 × 10 ⁻⁵
		0.02 5.7 × 10 ⁻⁴	4.685	
land_ice_flux_rms_tidal_velocity	0.05	0.01	2.806	0.133
		0.1	3.983	
land_ice_flux_topdragcoeff	0.0025	0.001	2.488	4.48 × 10 ⁻³
		0.01	4.637	
cvmix_kpp_criticalbulkrichardsonnumber	0.25	0.2	3.306	0.0558
		1.0	2.895	
cvmix_kpp_use_enhanced_diff	true (1.0)	false (0.0)	3.368	true (0.598)
cvmix_kpp_surface_layer_extent	0.1	0.07	3.173	0.0114
		0.14	3.353	
cvmix_background_diffusion	0.0	10-5	3.128	7.73 × 10 ⁻⁵
		10-4	2.470	
cvmix_background_viscosity	10-4	10-5	3.267	9.13 × 10 ⁻⁶
standardgm_tracer_kappa	1800	600	3.236	1620
		1200	3.440	
salinity_restoring_constant_piston_velocity	1.585 × 10 ⁻⁶	7 × 10 ⁻⁷	4.570	2.45 × 10 ⁻⁶
		3 × 10 ⁻⁶	4.106	





Ongoing work: Ice-ocean coupling in E3SM

Ingredients needed:

- Higher-order pressure gradient
 - thinner, more tilted layers
- Wetting-and-drying
 - Allows ice to advance and retreat






Higher-order pressure gradient

Accurate evaluation of contour integral:

$$\oint \rho(\mathbf{k}, \mathbf{l}) dl,$$

- Known approaches:
 - Choose EOS so integral is analytic (e.g. Adcroft et al. 2008)
 - high-order Gaussian quadrature
 (e.g. Engwirda et al. 2016)
- Should allow rx1 >> 5




From Shchepetkin and McWilliams (2003)





Wetting and Drying Scheme

- Thin film, O(cm) or less, of ocean in "dry" regions
- "Dry" cells allow mass fluxes in but prevent fluxed out
- Implemented for coastal sea-level change
- Work in progress: ice sheet advance/retreat:
 - Increased ice pressure from ice advance will cause drying
 - Decreased pressure from ice retreat will allow wetting




Example of ice-shelf retreat from a POPSICLES simulation





Summary

- E3SM v1.0 code released in April
 - low, medium, and high-resolution configurations included
 - Ice-shelf configurations included: use at your own risk
- Ice shelves will (hopefully) be default in E3SM v2.0
 - improve ocean properties, particularly at high resolution
 - allow projections of Antarctic melt rates and ultimately ice mass loss
- Dynamic coupling with land-ice model is ongoing
 - higher-order pressure gradient
 - wetting and drying



eddy activity in high-resolution ocean simulation



